arXiv:1207.1067v3 [math.NT] 16 Jul 2012

Bounding the Growth Rate of Approximation Coefficients using the Symmetry of Jager Pairs

Avraham Bourla
Department of Mathematics
Saint Mary's College of Maryland
Saint Mary's City, MD, 20686
abourla@smcm.edu

July 17, 2012

Abstract

Symmetrical subdivisions in the space of Jager Pairs will provide us with bounds on the rate of growth for the sequence of approximation coefficients corresponding to continued fractions like expansions. Results will also apply to the classical regular and backwards continued fractions expansions, which are realized as special cases.

1 Introduction

Given a real number r and a rational number, written as the unique quotient $\frac{p}{q}$ of the relatively prime integers p and $q > 0$, our fundamental object of interest from diophantine approximation is the **approximation coefficient** $\theta(r, \frac{p}{q}) := q^2\Big|r - \frac{p}{q}\Big|$. Small approximation coefficients correspond to high quality approximations, combining accuracy with simplicity. For instance, the error in approximating π using $\frac{355}{113} = 3.1415920353982$ is smaller than the error of its decimal expansion to the fifth digit $3.14159 = \frac{314159}{100000}$. Since the former rational also has a much smaller denominator, it is of far greater quality than the latter. Indeed $\theta\big(\pi, \frac{355}{113}\big) < 0.0341$ whereas $\theta\big(\pi, \frac{314159}{100000}\big) > 26,535$.

Adding any integer to a fraction does not change its denominator, hence we may restrict our attention to the unit interval. More specifically, we have $\theta(r, \frac{p}{q}) = \theta(x, \frac{p}{q} - \lfloor r \rfloor)$, where $\lfloor r \rfloor$ is the largest integer smaller than or equal to r (a.k.a. the the floor of r) and $x := r - \lfloor r \rfloor \in [0, 1)$. Expanding an irrational initial seed $x_0 \in (0, 1) - \mathbb{Q}$ as an infinite regular continued fraction

$$x_0 = [a_1, a_2, a_3, ...]_0 := \cfrac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + ...}}}$$

provides us with the unique symbolic representation via the sequence $\{a_n\}_1^\infty$ of positive integers, known as the partial quotients or **digits of expansion**. For all $n \geq 1$, we label the approximation coefficient associated with the n$^{\text{th}}$ **convergent** of x_0, $\frac{p_n}{q_n} := [a_1.a_2, ..., a_n]_0$ by θ_n and refer to the sequence $\{\theta_n\}_1^\infty$ as the **sequence of approximation coefficients**.

For all $n \geq 1$, it is well known [3, Theorem 4.6] that $\left|x_0 - \frac{p_n}{q_n}\right| < \frac{1}{q_n q_{n+1}} < \frac{1}{q_n^2}$, hence $\theta_n < 1$. Conversely, Legendre [3, Theorem 5.12] proved that if $\theta(x_0, \frac{p}{q}) < \frac{1}{2}$ then $\frac{p}{q}$ is a convergent of x_0. In 1891, Hurwitz [5, Theorem 5.1.4] showed there exist infinitely many pairs of integers p and q such that the inequality $\theta\left(x_0, \frac{p}{q}\right) < \frac{1}{\sqrt{5}}$ is sharp. Conclude that all irrational numbers possess infinitely many high quality approximations using rational numbers, whose associated approximation coefficients are less than $\frac{1}{\sqrt{5}}$, all of which must belong to the sequence of regular continued fraction convergents for x_0. In fact, the value of $\liminf_{n\to\infty}\{\theta_n(x_0)\}$ measures how well can x_0 be approximated by rational numbers, leading to the construction of the Lagrange Spectrum (for more details, refer to [4]). In 1978, Jurkat and Peyerimhoff [8] showed that the **Space of Jager Pairs**

$$\left\{(\theta_n(x_0), \theta_{n+1}(x_0)) \subset \mathbb{R}^2, \, x_0 \in (0,1) - \mathbb{Q}, \, n \geq 1\right\}$$

is a dense subset of the interior of the triangle in the cartesian plane with vertices $(0,0)$, $(0,1)$ and $(1,0)$.

Our goal is two folded: we will first show that the symmetry in this space discovered in 2002 by Dajani and Kraaikamp [5], extends to the two one parameter families of continued fractions-like expansions, first introduced in 2003 by Haas and Molnar [6]. In particular, the results will apply to the Space of Jager Pairs associated with the backwards continued fractions expansion

$$x_0 = 1 - \cfrac{1}{a_1 + 1 - \cfrac{1}{a_2 + 1 - \cfrac{1}{a_3 + \ldots}}}.$$

Next, a simple plane geometry argument will provide us with upper and lower bounds for the growth rate of the sequence of approximation coefficients, given some a-priori knowledge of the digits of expansion. For instance, if $a_2 = a_3 = 1$ and $a_4 = 3$ are the digits of the classical regular continued fraction expansion, then corollary 4.4 will imply that $|\theta_2 - \theta_1| < \frac{\sqrt{2}}{3}$ and $\frac{2\sqrt{2}}{7} < |\theta_3 - \theta_2| < \frac{3\sqrt{2}}{5}$.

2 Preliminaries

This section is a paraphrased summery of excerpts from [6, 7], given for sake of completeness. In general, the fractional part of Möbius transformation which map $[0,1]$ onto $[0,\infty]$ leads to expansion of real numbers as continued fractions. To characterize all these transformations, we recall that Möbius transformations are uniquely determined by their values for three distinct points. Thus, we will need to introduce a parameter for the image of an additional point besides 0 and 1, which we will naturally take to be ∞. Since our maps fix the real line, the image of ∞, denoted by $-k$, can take any value within the set of negative real numbers. We let $m \in \{0,1\}$ equal zero for orientation reversing transformations, i.e. $0 \mapsto \infty$, $1 \mapsto 0$ and let m equal one for orientation preserving transformations, i.e. $0 \mapsto 0$, $1 \mapsto \infty$. Conclude that all such transformations must be the extension of the maps $x \mapsto \frac{k(1-m-x)}{x-m}$, $k > 0$, mapping $(0,1)$ homeomorphically to $(0,\infty)$, to the extended complex plane. The maps $T_{(m,k)} : [0,1) \to [0,1)$, $0 \mapsto 0$,

$$T_{(m,k)}(x) = \frac{k(1-m-x)}{x-m} - \left\lfloor \frac{k(1-m-x)}{x-m} \right\rfloor, \quad x > 0$$

are called **Gauss-like** and **Renyi-like** for $m = 0$ and $m = 1$ respectively.

We expand the initial seed $x_0 \in (0,1)$ as an (m,k)-continued fraction using the following iteration process:

1. Set $n := 1$.
2. If $x_{n-1} = 0$, write the (m,k)-expansion of x_0 as $[a_1, ..., a_{n-1}]_{(m,k)}$ and exit.
3. Set the **digit** and **future** of x_0 at time n to be $a_n := \left\lfloor \frac{k(1-m-x_{n-1})}{x_{n-1}-m} \right\rfloor \in \mathbb{Z}^+ := \mathbb{Z} \cap [0,\infty)$ and $x_n := \frac{k(1-m-x_{n-1})}{x_{n-1}-m} - a_n \in [0,1)$. Increase n by one and goto step 2.

For all $n \geq 0$, we thus have

$$x_{n+1} = T_{(m,k)}(x_n) = \frac{k(1-m-x_n)}{x_n - m} - a_n$$

so that

$$x_n = m + \frac{k(1-2m)}{a_{n+1} + k + x_{n+1}}.$$

Therefore, this iteration scheme leads to the expansion of the initial seed x_0 as

$$x_0 = m + \frac{k(1-2m)}{a_1 + k + x_1} = m + \frac{k(1-2m)}{a_1 + k + m + \frac{k(1-2m)}{a_2+k+x_2}} = ...$$

Remark 2.1. The classical $k = 1$ cases lead to the regular and backwards continued fractions expansions for $m = 0$ and $m = 1$ respectively, bit with digits which are smaller by one than their classical representation. For instance,

$$[0,1,2]_{(0,k)} = \cfrac{k}{0 + k + \cfrac{k}{1 + k + \cfrac{k}{2+k}}} = \frac{k^2+4k+2}{k^2+5k+4}$$

and

$$[0,1,2]_{(1,k)} = 1 - \cfrac{k}{0 + k + 1 - \cfrac{k}{1 + k + 1 - \cfrac{k}{2+k}}} = \frac{k+4}{k^3+3k^2+5k+4}$$

will yield, after plugging $k = 1$, the fractions $[1,2,3]_0 = \frac{7}{10}$ and $[1,2,3]_1 = \frac{5}{13}$.

Since for the classical regular and backwards continued fraction expansions, this iteration process eventually terminates precisely when the initial seed x_0 is a rational number. Analogously, we denote the countable set of all numbers in the interval with finite (m,k)-expansion by $\mathbb{Q}_{(m,k)}$. For all $a \in \mathbb{Z}^+$, the cylinder set

$$\Delta_a := \left(\frac{(1-m)k+ma}{a+k+1-m}, \frac{(1-m)k+m(a+1)}{a+k+m}\right) \tag{1}$$

is defined such that $x_0 \in \Delta_a$ iff $a_1 = a$ or, more generally,

$$x_n \in \Delta_a \iff a_{n+1} = a, \quad n \geq 0. \tag{2}$$

The restriction of the map $T_{(m,k)}$ to Δ_a is a homeomorphism onto (0,1). We further define the **past** of x_0 at time $n \geq 2$ to be

$$Y_n(x_0) = Y_n := m-k-a_n-[a_{n-1},...,a_1]_{(m,k)} \in (m-k-a_n-1, m-k-a_n). \tag{3}$$

The set $\big\{(x_n, Y_n) : x_0 \in (0,1) - \mathbb{Q}_{(m,k)}, n \geq 2\big\}$ is a subset of the space $\Omega_{(m,k)} := (0,1) \times (-\infty, m-k)$.

The sequence of approximation coefficients $\{\theta_n(x_0)\}_1^\infty$ for the (m,k)-expansion is defined just like the classical object, i.e. $\theta_n(x_0) := q_n^2\left|x_0 - \frac{p_n}{q_n}\right|$, where $\frac{p_n}{q_n} = [a_1,...,a_n]_{(m,k)} \in \mathbb{Q}_{(m,k)}$ are the appropriate convergents. The sequence of approximation coefficients relates to the future and past sequences of x_0 using the identity

$$\theta_n(x_0) = \frac{1}{x_{n+1} - Y_{n+1}},$$

first proven for the classical regular continued fraction case in 1921 by Perron [9]). The equicontinuous family of continuous maps

$$\Psi_{(m,k)} : \Omega_{(m,k)} \to \mathbb{R}^2, \quad (x,y) \mapsto \left(\frac{1}{x-y}, \frac{(m-x)(m-y)}{(2m-1)k(x-y)}\right), \tag{4}$$

$$(\theta_n, \theta_{n+1}) = \Psi_{(m,k)}(x_{n+1}, Y_{n+1}).$$

Each map is a homeomorphism onto its image $\Psi_{(m,k)}(\Omega_{(m,k)})$, which we will denote by $\Gamma_{(m,k)}$. the corresponding **Space of Jager Pairs**

$$\{(\theta_n(x_0), \theta_{n+1}(x_0)) : x_0 \in (0,1) - \mathbb{Q}_{(m,k)}, n \geq 1\}$$

is a dense subset of $\Gamma_{(m,k)}$. In our excursion, we will require the parameter k to be grater than one, treating the classical regular and backwards continued fraction expansions as the limit of the (m,k)-expansions. As $k \to 1^+$. The choice of taking the limit from the right is not arbitrary, for the treatment of the $0 < k < 1$ cases will exhibit certain pathologies (for instance, see [2, section 3.3]).

3 The finer structure for the Space of Jager Pairs

Fix $m \in \{0,1\}$, $k \in (1,\infty)$, and an initial seed $x_0 \in (0,1) - \mathbb{Q}_{(m,k)}$. In order to ease the notation, we will omit the subscripts $\square_{(m,k)}$ from now on. For all $a \in \mathbb{Z}^+$, define define the regions

$$P_{(m,k,a)} = P_a := (0,1) \times (m-a-k-1, m-a-k)$$

and

$$F_{(m,k,a)} = F_a := \Delta_a \times (-\infty, m-k),$$

where Δ_a is the cylinder set (1). Using the identity (2) and the definition (3) of Y_n, we see that that for all $n \geq 1$ we have

$$(x_{n+1}, y_{n+1}) \in P_a \cap F_b \iff a_{n+1} = a \text{ and } a_{n+2} = b.$$

We label the images of each of these subsets of Ω under Ψ by $P_a^\#$ and $F_a^\#$. Using formula (4) and the fact that Ψ is a homeomorphism, we see that for all $n \geq 1$, we have

$$(\theta_n, \theta_{n+1}) \in P_a^\# \cap F_b^\# \iff a_{n+1} = a \text{ and } a_{n+2} = b. \tag{5}$$

Next, let $p_{(m,k,a)} = p_a$ be the open horizontal line segment $(0,1)\times\{m-a-k\}$, let $f_{(m,k,a)} = f_a$ be the open vertical ray $\left\{\frac{(1-m)k+m}{a+k}\right\} \times (-\infty, m-k)$ and let $p_a^\#$ and $f_a^\#$ be their image under Ψ. Since both the collections $\{P_a \cup p_a\}_{a\in\mathbb{Z}^+}$ and $\{F_a \cup f_a\}_{a\in\mathbb{Z}^+}$ partition Ω, the image of their intersections under Ψ, $\left\{(P_a^\# \cup p_a^\#) \cap (F_b^\# \cup f_b^\#)\right\}_{a,b\in\mathbb{Z}^+}$ will partition Γ. We will call each member of this refined partition a **subdivision**.

Proposition 3.1. For all $k \in \mathbb{R} > 1$ and $a, b \in \mathbb{Z}^+$ the region $P_a^\# \cap F_b^\#$ is the open interior of the quadrangle in $\mathbb{E}^2$ with vertices

$$\left(\frac{b+k}{(1-2m)k+(a+k)(b+k)}, \frac{a+k}{(1-2m)k+(a+k)(b+k)}\right),$$

$$\left(\frac{b+k}{(1-2m)k+(a+k+1)(b+k)}, \frac{a+k+1}{(1-2m)k+(a+k+1)(b+k)}\right),$$

$$\left(\frac{b+k+1}{(1-2m)k+(a+k)(b+k+1)}, \frac{a+k}{(1-2m)k+(a+k)(b+k+1)}\right)$$

and

$$\left(\frac{b+k+1}{(1-2m)k+(a+k+1)(b+k+1)}, \frac{a+k+1}{(1-2m)k+(a+k+1)(b+k+1)}\right).$$

Proof. Fix $a \in \mathbb{Z}^+$ and for every $(x, y_0) \in p_a$ i.e. $x \in (0,1)$, $y_0 := m-k-a$, set $(u,v) := \Psi(x, y_0)$. Using the definition (4) of Ψ, we write

$$u = \frac{1}{x-y_0} = \frac{1}{x+a+k-m} : \frac{1}{a+k-m} \to \frac{1}{a+k+1-m} \quad \text{as } x : 0 \to 1.$$

Since $m - x = m - y - \frac{1}{u}$, we express v in terms of u as

$$v = \frac{(m-x)(m-y_0)}{(2m-1)k(x-y_0)} = \frac{(m-x)(m-y_0)u}{(2m-1)k} = \frac{a+k}{(2m-1)k}\left((a+k)u-1\right)$$

so that

$$v : \frac{m(a+k)}{k(a+k-m)} \to \frac{(1-m)(k+a)}{k(k+a+1-m)} \quad \text{as } x : 0 \to 1.$$

Since $\Psi : \Omega \to \Gamma$ is a homeomorphism, we see that $p_a^{\#}$ is an open segment of the line $(a+k)^2u+(1-2m)kv = a+k$ between the points $\left(\frac{1}{a+k-m}, \frac{m(a+k)}{k(a+k-m)}\right)$ and $\left(\frac{1}{a+k+1-m}, \frac{(1-m)(k+a)}{k(a+k+1-m)}\right)$.


$\frac{1}{k}$
$\left(\frac{1}{k+4}, \frac{k+3}{k(k+4)}\right)$
$\left(\frac{1}{k+3}, \frac{k+2}{k(k+3)}\right)$
$\left(\frac{1}{k+2}, \frac{k+1}{k(k+2)}\right)$
$\left(\frac{1}{k+1}, \frac{1}{k+1}\right)$
$P_2^{\#}$
$P_1^{\#}$
$P_0^{\#}$
$\frac{1}{k+3}$
$\frac{1}{k+2}$
$\frac{1}{k+1}$
$\frac{1}{k}$

$\Gamma_{(0,k)}$


$\left(\frac{1}{k-1}, \frac{1}{k-1}\right)$
$\left(\frac{1}{k}, \frac{k+1}{k^2}\right)$
$\left(\frac{1}{k+1}, \frac{k+2}{k(k+1)}\right)$
$\left(\frac{1}{k+2}, \frac{k+3}{k(k+2)}\right)$
$P_2^{\#}$
$P_1^{\#}$
$P_0^{\#}$
$\frac{1}{k+3}$
$\frac{1}{k+2}$
$\frac{1}{k+1}$
$\frac{1}{k}$

$\Gamma_{(1,k)}$

Next, fix $b \in \mathbb{Z}^+$ and set $x_0 := \frac{(1-m)k+mb}{b+k}$. We use the definition (4) of Ψ to first write $m - y = m - x + \frac{1}{u}$ and then express v in terms of u as

$$v = \frac{u}{(2m-1)k}(m-x_0)\left(m - x_0 + \frac{1}{u}\right) = \frac{1}{b+k}\left(\frac{(2m-1)k}{b+k}u + 1\right). \tag{6}$$

Since Ψ is a homeomorphism, this allows us to conclude that Ψ maps f_b to an open segment of a line in the uv plane, which is the reflection of the line $p_b^{\#}$ along the diagonal $u - v = 0$. In particular, $P_a^{\#} \cap F_b^{\#}$ is the region interior to the quadrangle with vertices $p_a^{\#} \cap f_b^{\#}$, $p_{a+1}^{\#} \cap f_b^{\#}$, $p_a^{\#} \cap f_{b+1}^{\#}$ and $p_{a+1}^{\#} \cap f_{b+1}^{\#}$. But $(x_0, y_0) = p_a \cap f_b$, so that $(u_0, v_0) := \Psi(x_0, y_0) = p_a^{\#} \cap f_b^{\#}$. The definition (4) of Ψ and formula (6) now yield

$$u_0 = \frac{1}{x_0 - y_0} = \frac{b+k}{(1-2m)k + (a+k)(b+k)}$$

and

$$v_0 = \frac{a+k}{(1-2m)k + (a+k)(b+k)}$$

as desired. ■

When $m = 0$, we obtain the depicted Spaces of Jager Pairs:


$\frac{1}{k}$
$\left(\frac{1}{k+4}, \frac{k+3}{k(k+4)}\right)$
$\left(\frac{1}{k+3}, \frac{k+2}{k(k+3)}\right)$
$\left(\frac{1}{k+2}, \frac{k+1}{k(k+2)}\right)$
$\left(\frac{1}{k+1}, \frac{1}{k+1}\right)$
$\frac{1}{k+1}$
$\left(\frac{k+1}{k(k+2)}, \frac{1}{k+2}\right)$
$\frac{1}{k+2}$
$\left(\frac{k+2}{k(k+3)}, \frac{1}{k+3}\right)$
$\frac{1}{k+3}$
$\left(\frac{k+3}{k(k+4)}, \frac{1}{k+4}\right)$
$P_2^\# \cap F_0^\#$
$P_1^\# \cap F_0^\#$
$P_0^\# \cap F_0^\#$
$P_2^\# \cap F_1^\#$
$P_1^\# \cap F_1^\#$
$P_0^\# \cap F_1^\#$
$P_2^\# \cap F_2^\#$
$P_1^\# \cap F_2^\#$
$P_0^\# \cap F_2^\#$
$\frac{1}{k+3}$
$\frac{1}{k+2}$
$\frac{1}{k+1}$
$\frac{1}{k}$

$\Gamma_{(0,k)}$ revisited


(0, 1)
$\left(\frac{1}{5}, \frac{4}{5}\right)$
$\left(\frac{1}{4}, \frac{3}{4}\right)$
$\left(\frac{1}{3}, \frac{2}{3}\right)$
$\left(0, \frac{1}{2}\right)$
$\left(\frac{2}{3}, \frac{1}{3}\right)$
$\left(0, \frac{1}{3}\right)$
$\left(\frac{3}{4}, \frac{1}{4}\right)$
$\left(0, \frac{1}{4}\right)$
$\left(\frac{4}{5}, \frac{1}{5}\right)$
$P_2^\# \cap F_0^\#$
$P_1^\# \cap F_0^\#$
$P_0^\# \cap F_0^\#$
$P_2^\# \cap F_1^\#$
$P_1^\# \cap F_1^\#$
$P_0^\# \cap F_1^\#$
$P_2^\# \cap F_2^\#$
$P_1^\# \cap F_2^\#$
$P_0^\# \cap F_2^\#$
$\left(\frac{1}{4}, 0\right)$
$\left(\frac{1}{3}, 0\right)$
$\left(\frac{1}{2}, 0\right)$
(1, 0)

$\Gamma_{(0,1)}$

This result was first proved for the classical regular continued fractions corresponding to $k = 1$ by Dajani and Kraaikamp [5, exercise 5.3.4]. For this case, the region $P_0^\# \cap F_0^\#$ degenerates to the interior of the triangle with vertices $\left(\frac{1}{3}, \frac{2}{3}\right)$, $\left(\frac{2}{3}, \frac{1}{3}\right)$ and $\left(\frac{2}{5}, \frac{2}{5}\right)$. When $m = 1$, we obtain the depicted Spaces of Jager Pairs:


$\left(\frac{1}{k-1}, \frac{1}{k-1}\right)$
$\left(\frac{1}{k}, \frac{k+1}{k^2}\right)$
$\left(\frac{1}{k+1}, \frac{k+2}{k(k+1)}\right)$
$\left(\frac{1}{k+2}, \frac{k+3}{k(k+2)}\right)$
$\frac{1}{k}$
$\left(\frac{k+1}{k^2}, \frac{1}{k}\right)$
$\frac{1}{k+1}$
$\left(\frac{k+2}{k(k+1)}, \frac{1}{k+1}\right)$
$\frac{1}{k+2}$
$\left(\frac{1}{k+2}, \frac{k+3}{k(k+2)}\right)$
$\frac{1}{k+3}$
$P_2^\# \cap F_0^\#$
$P_1^\# \cap F_0^\#$
$P_0^\# \cap F_0^\#$
$P_2^\# \cap F_1^\#$
$P_1^\# \cap F_1^\#$
$P_0^\# \cap F_1^\#$
$P_2^\# \cap F_2^\#$
$P_1^\# \cap F_2^\#$
$P_0^\# \cap F_2^\#$
$\frac{1}{k+3}$
$\frac{1}{k+2}$
$\frac{1}{k+1}$
$\frac{1}{k}$

$\Gamma_{(1,k)}$ revisited



$\Gamma_{(1,1)}$

The picture for the classical backwards continued fractions $k = 1$, is obtained after we let $k \to 1^+$ and use the equicontinuity of the family $\{\Psi_{(1,k)}\}_{k>1}$. For this case, the region $P_0^\# \cap F_0^\#$ expands to the unbounded region in the uv plane, which is the intersection of the regions $u - v > 1$, $v - u < 1$, $4u - v > 2$ and $4v - u > 2$.

4 Results

This symmetry exhibited by the subdivisions in the Space of Jager Pairs will allow us to provide uniform bounds for the **growth rate sequence** for $\{\theta_n\}_1^\infty$, which is the sequence whose member at time n is $\{|\theta_{n+1} - \theta_n|\}$. But first, we already know that $\Gamma_{(1,1)}$ is the unbounded region in the first quadrant of the uv plane bounded by the lines $u - v < 1$ and $v - u < 1$, hence we obtain at once that

Theorem 4.1. In general, the sequence of approximation coefficients associated with the classical backwards continued fraction expansion $m = k = 1$ has no uniform upper bound. However its growth rate sequence is uniformly bounded by one.

Our main result is:

Theorem 4.2. Given $m \in \{0, 1\}$, $k \in \mathbb{R} > 1$, $x_0 \in (0, 1) - \mathbb{Q}_{(m,k)}$ and $N \geq 1$, write $l := \min\{a_{N+1}, a_{N+2}, a_{N+3}\}$ and $L := \max\{a_{N+1}, a_{N+2}, a_{N+3}\}$, where a_n is the digit at time n in the (m,k)-expansion for x_0. Then the inequality

$$(\theta_{N+1} - \theta_N)^2 + (\theta_{N+2} - \theta_{N+1})^2 < 2\left(\frac{L - l + 1}{(1 - 2m)k + (l + k)(L + k + 1)}\right)^2$$

is sharp. Furthermore, if $L - l > 1$, then the inequality

$$(\theta_{N+1} - \theta_N)^2 + (\theta_{N+2} - \theta_{N+1})^2 > 2\left(\frac{L - l}{(1 - 2m)k + (l + k + 1)(L + k)}\right)^2$$

is sharp.

To prove this theorem, we will first prove:

Lemma 4.3. Assuming the hypothesis of the theorem, write $a := \min\{a_{N+1}, a_{N+2}\}$, $A := \max\{a_{N+1}, a_{N+2}\}$, $b := \min\{a_{N+2}, a_{N+3}\}$ and $B := \max\{a_{N+2}, a_{N+3}\}$. Then the inequality

$$\begin{aligned}&(\theta_{N+1} - \theta_N)^2 + (\theta_{N+2} - \theta_{N+1})^2 \\ &< \left(\frac{B + k + 1}{(1 - 2m)k + (a + k)(B + k + 1)} - \frac{b + k}{(1 - 2m)k + (b + k)(A + k + 1)}\right)^2 \\ &+ \left(\frac{A + k + 1}{(1 - 2m)k + (b + k)(A + k + 1)} - \frac{a + k}{(1 - 2m)k + (a + k)(B + k + 1)}\right)^2\end{aligned}$$

is sharp. Furthermore, if $\max\{A - a, B - b > 1\}$, then the inequality

$$(\theta_{N+1} - \theta_N)^2 + (\theta_{N+2} - \theta_{N+1})^2$$

$$> \left(\frac{B+k}{(1-2m)k+(a+k+1)(B+k)} - \frac{b+k+1}{(1-2m)k+(b+k+1)(A+k)}\right)^2$$
$$+ \left(\frac{A+k}{(1-2m)k+(b+k+1)(A+k)} - \frac{a+k+1}{(1-2m)k+(a+k+1)(B+k)}\right)^2$$

is sharp.

Proof. Let $R_{(m,k,a,b,A,B)} = R$ be the set

$$\bigcup_{a\le i\le A,\, b\le j\le B} (P_i^{\#} \cap F_j^{\#}),$$

so that, from formula (5), we obtain that $(\theta_N, \theta_{N+1}) \in P^{\#}_{a_N+1} \cap F^{\#}_{a_{N+2}} \subset R$ as well as $(\theta_{N+1}, \theta_{N+2}) \in P^{\#}_{a_{N+2}} \cap F^{\#}_{a_{N+3}} \subset R$. From proposition 3.1, we see that R is the interior of the bounded convex quadrangle, consisting of sections from the line segments $p_a^{\#}$, $p_{A+1}^{\#}$, $f_b^{\#}$ and $f_{B+1}^{\#}$. Unless $\max\{A-a, B-b\} \le 1$, we also let $r_{(m,k,a,b,A,B)} = r$ be the non-empty set

$$\bigcup_{a+1\le i\le A-1,\, b+1\le j\le B-1} (P_i^{\#} \cap F_j^{\#}),$$

so that

$$\{(\theta_N, \theta_{N+1}), (\theta_{N+1}, \theta_{N+2})\} \subset R - r. \tag{7}$$

From the same proposition, we see that $r \subset R$ is the interior of the convex quadrangle, consisting of sections from the segments $p_{a+1}^{\#}$, $p_A^{\#}$, $f_{b+1}^{\#}$ and $f_B^{\#}$, that is, the quadrangle r is obtained from peeling the outer layer of R's subdivisions.

When $m = 0$, let $(u_0, v_0) := p_A^{\#} \cap f_{b+1}^{\#}$ and $(U_0, V_0) := p_{a+1}^{\#} \cap f_B^{\#}$. Then

$$(u, v) \in P_A^{\#} \cap F_b^{\#} \implies u < u_0 \text{ and } v > v_0$$

and

$$(u, v) \in P_a^{\#} \cap F_B^{\#} \implies u > U_0 \text{ and } v < V_0.$$

When $m = 1$, let $(u_1, v_1) := p_A^{\#} \cap f_B^{\#}$ and $(U_1, V_1) := p_{a+1}^{\#} \cap f_{b+1}^{\#}$. Then

$$(u, v) \in P_a^{\#} \cap F_b^{\#} \implies u < u_0 \text{ and } v < v_0$$

and

$$(u, v) \in P_A^{\#} \cap F_B^{\#} \implies u > U_0 \text{ and } v > V_0.$$

In tandem with formula (7), this allows us to conclude that for either case we have

$$\operatorname{diam}(r)^2 < d\left((\theta_N, \theta_{N+1}), (\theta_{N+1}, \theta_{N+2})\right) = (\theta_N - \theta_{N+1})^2 + (\theta_{N+1} - \theta_{N+2})^2 < \operatorname{diam}(R)^2, \tag{8}$$

where d is the familiar distance formula between two points in $\mathbb{E}^2$ and $\operatorname{diam}(R)$ stands for the euclidean diameter of the region R.

The diameter of the interior of a convex quadrangle in $\mathbb{E}^2$ with opposite acute angles is the length of the diagonal connecting the vertices corresponding to these acute angles. To verify this simple observation from plane geometry, observe that from the continuity of the euclidean distance formula, the diameter of a convex polygon is either a side or a diagonal. The diagonal connecting the two acute angles will lie opposite to an obtuse angle in either one of the induced triangles obtained. Since for any triangle, the length of a side is proportional to the measure of its opposite angle, we conclude that this diagonal is longer than all of the four sides in this quadrangle. To verify it is longer than the other diagonal, consider one of the four induced triangles obtained after drawing both diagonals and use the same opposite angle argument.

When $m = 0$ and $c \in \mathbb{Z}^+$, the slope of the line containing $p_c^\#$ lie in $(-\infty, -1)$ whereas the slope of the line containing $f_c^\#$ lie in $(-1, 0)$. Conclude that both R and r have opposite acute angles, so that their diameters are the length of the diagonal connecting the vertices $p_a^\# \cap f_{B+1}^\#$ with $p_{A+1}^\# \cap f_b^\#$ and $p_{a+1}^\# \cap f_B^\#$ with $p_A^\# \cap f_{b+1}^\#$ respectively. Hence

$$\mathrm{diam(R)} = \mathrm{d}(p_a^\# \cap f_{B+1}^\#, p_{A+1}^\# \cap f_b^\#)$$

and

$$\mathrm{diam(r)} = \mathrm{d}(p_{a+1}^\# \cap f_B^\#, p_A^\# \cap f_{b+1}^\#).$$

When $m = 1$, the slope of the line containing $p_c^\#$ lie in $(1, \infty)$ whereas the slope of the line containing $f_c^\#$ lie in $(0, 1)$. Conclude that both R and r has opposite acute angles, so that their diameters are the length of the diagonal connecting the vertices $p_a^\# \cap f_b^\#$ with $p_{A+1}^\# \cap f_{B+1}^\#$ and $p_{a+1}^\# \cap f_{b+1}^\#$ with $p_A^\# \cap f_B^\#$ respectively. Hence

$$\mathrm{diam(R)} = \mathrm{d}(p_a^\# \cap f_b^\#, p_{A+1}^\# \cap f_{B+1}^\#)$$

and

$$\mathrm{diam(r)} = \mathrm{d}(p_{a+1}^\# \cap f_{b+1}^\#, p_A^\# \cap f_B^\#).$$

For either case, we have

$$\mathrm{diam(R)}^2 = \left(\frac{B+k+1}{(1-2m)k + (A+k+1)(B+k+1)} - \frac{b+k}{(1-2m)k + (a+k)(b+k)}\right)^2 + \left(\frac{A+k+1}{(1-2m)k + (A+k+1)(B+k+1)} - \frac{a+k}{(1-2m)k + (a+k)(b+k)}\right)^2$$

and

$$\mathrm{diam(r)}^2 = \left(\frac{B+k}{(1-2m)k + (A+k)(B+k)} - \frac{b+k+1}{(1-2m)k + (a+k+1)(b+k+1)}\right)^2 + \left(\frac{A+k}{(1-2m)k + (A+k)(B+k)} - \frac{a+k+1}{(1-2m)k + (a+k+1)(b+k+1)}\right)^2.$$

Formula (8) now yields the desired inequalities and the density of the space of Jager Pairs in $\Gamma_{(m,k)}$ establishes their sharpness.

■

Proof. (of theorem 4.2). For all integers a, b, A and B with $l \leq a \leq A \leq L$ and $l \leq b \leq B \leq L$, we have

$$\frac{A+k+1}{(1-2m)k+(b+k)(A+k+1)} \leq \frac{L+k+1}{(1-2m)k+(l+k)(L+k+1)}$$

and

$$\frac{l+k}{(1-2m)k+(l+k)(L+k+1)} \leq \frac{a+k}{(1-2m)k+(a+k)(B+k+1)},$$

so that

$$\begin{aligned} 0 &\leq \frac{L-l}{(1-2m)k+(l+k+1)(L+k)} \\ &\leq \frac{A+k+1}{(1-2m)k+(b+k)(A+k+1)} - \frac{a+k}{(1-2m)k+(a+k)(B+k+1)} \\ &\leq \frac{L+k+1}{(1-2m)k+(l+k)(L+k+1)} - \frac{l+k}{(1-2m)k+(l+k)(L+k+1)} \\ &= \frac{L-l+1}{(1-2m)k+(l+k)(L+k+1)}. \end{aligned}$$

This argument remains identical after we exchange a for b and A for B. Furthermore, setting $a = b := l$ and $A = B := L$ shows that we cannot replace these weak inequalities with strict ones. The lemma will now provide us with the result. ■

Corollary 4.4. Assuming the hypothesis of the theorem, we have

$$\max\{|\theta_{N+1} - \theta_N|, |\theta_{N+2} - \theta_{N+1}|\} < \sqrt{2}\left(\frac{L-l+1}{(1-2m)k+(l+k)(L+k+1)}\right).$$

Furthermore, if $L - l > 1$, we also have

$$\min\{|\theta_{N+1} - \theta_N|, |\theta_{N+2} - \theta_{N+1}|\} > \sqrt{2}\left(\frac{L-l}{(1-2m)k+(l+k+1)(L+k)}\right).$$

These results extend to the classical $k = 1$ cases with one exception: the upper bound statement does not apply for the $m = 1$ case when $a_{n+2} = 0$ and either $a_{n+1} = 0$ or $a_{n+3} = 0$.

Proof. When $k > 1$ or when $k = 1$ and $m = 0$, this is an immediate consequence of the theorem. When $m = 1$, we let $k \to 1^+$ and use the equicontinuity of the family $\{\Psi_{(1,k)}\}_{k>1}$ to obtain the result, once we exclude the possibility of the unbounded region $P_0^{\#} \cap F_0^{\#}$ from belonging to the region R, when computing the upper bound. This happens precisely when either $a_{n+1} = a_{n+2} = 0$ or $a_{n+2} = a_{n+3} = 0$. ■

Remark 4.5. Since our digits are lower by one than their classical representation (see remark 2.1), the bounds in the example ending the introduction are obtained by applying this corollary to the digits $a_2 = a_3 = 0$ and $a_4 = 2$ in the (0,1)-expansion of the initial seed.

5 Acknowledgments

This paper is a development of part of the author's Ph.D. dissertation at the University of Connecticut. The author has benefited tremendously from the patience and rigor his advisor Andy Haas and would like to thank him for all his efforts. He would also like to extend his gratitude to Cor Kraaikamp and Matt Papanikolas, whose suggestions had contributed to a more clear and elegant finished product.

References

[1] -, *Symmetry in the sequence of approximation coefficients*, Proceedings of the Amer. Math. Soc., to appear.

[2] -, *The bi-sequence of approximation coefficients for Gauss-like and Renyi-like maps on the interval*, Ph.D. Thesis, University of Connecticut press, 2011.

[3] E. B. Burger, *Exploring the number jungle: A journey into diophantine analysis*, Providence, RI: Amer. Math. Soc., 2000.

[4] T. W. Cusick and M. E. Flahive, *The Markoff and Lagrange spectra*, Mathematical Surveys and Monograms no. 30, Amer. Math. Soc., 1989.

[5] K. Dajani and C. Kraaikamp, *Ergodic theory of numbers*, The Carus Mathematical Monograms no. 29, Math. Asso .Amer. , 2002.

[6] A. Haas and D. Molnar, *Metrical diophantine approximation for continued fractions like maps of the interval*, Transaction of the Amer. Math. Soc., Volume 356(7), 2003.

[7] A. Haas and D. Molnar, *The distribution of Jager pairs for continued fraction like mappings of the interval*, Pacific J. Math. 217(1), 2004.

[8] W. B. Jurkat and A. Peyerimhoff, *Characteristic approximation properties of quadratic irrationals*, Internat. J. Math. & Math. Sci.(1), 1978.

[9] O. Perron, *Über die approximation irrationaler zahlen durch rationale*, Heindelberg Akad. Wiss. Abh.(4), 1921.